

For your college-bound patients with partial-onset seizures, going back to school can pose many challenges. Now is a good time to help prepare them for the upcoming school year.

Being **Proactive** and **Simplifying** their regimen by reducing their number of daily doses can make a difference!

Make the switch to once-daily MOTPOLY XR

The established efficacy and tolerability of lacosamide... now in a **single daily dose**



MOTPOLY XR comes as a 100-mg, 150-mg, and 200-mg capsule

INDICATION

MOTPOLY XR is indicated for the treatment of partial-onset seizures in adults and in pediatric patients weighing at least 50 kg.

IMPORTANT SAFETY INFORMATION WARNINGS AND PRECAUTIONS

- Antiepileptic drugs increase the risk of suicidal behavior and ideation. Monitor patients for the emergence or worsening of depression, suicidal thoughts or behaviors.
- MOTPOLY XR may cause dizziness and ataxia in patients. Advise patients not to operate machinery or motor vehicles until they know how MOTPOLY XR affects them.
- Obtain ECG before beginning MOTPOLY XR, and after titration to steady-state maintenance dose in patients with underlying proarrhythmic conditions or those on concomitant medications that affect cardiac conduction. Closely monitor these patients.
- MOTPOLY XR may cause syncope in patients.

Please see additional Important Safety Information on reverse side.



Seizures School

For patients weighing ≥50 kg with partial-onset seizures

ONE AND DONE





For a college student with partial-onset seizures, like Shawn, consider once-daily MOTPOLY XR

The established efficacy and tolerability of lacosamide...now in a single daily dose1

- Steady 24-hour delivery with fewer fluctuations
- Proven XR bead technology achieves once-daily dosing
- Bioequivalent to lacosamide at steady-state blood levels

Educate Students and Their Parents on Tips for Back-to-School Success

Develop a support network²

- Become familiar with the college's health services
- Inform roomates and other friends about your epilepsy
- Inform the Resident Assistant (RA), support staff, and coaches

Take care of yourself³

- Get enough sleep as lack of sleep can be a seizure trigger
- Stay connected to friends and family

WARNINGS AND PRECAUTIONS (cont)

- Gradually withdraw MOTPOLY XR to minimize the potential of increased seizure frequency.
- Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)/multi-organ hypersensitivity has been reported and can be life-threatening or fatal. If signs or symptoms are present, immediately evaluate the patient. Discontinue MOTPOLY XR if there is no alternative etiology.

MOST COMMON ADVERSE REACTIONS

The most common adverse reactions in adults (≥10% and greater than placebo) are diplopia, headache, dizziness, nausea, and somnolence.

PREGNANCY: Based on animal data, MOTPOLY XR may cause fetal harm.

DRUG INTERACTIONS: Consider dose reduction in patients with renal or hepatic impairment taking strong inhibitors of CYP3A4 and CYP2C9.

Please see full Prescribing Information enclosed.

